

NOROVIRUS FACTSHEET

DISTRICT OF COLUMBIA DEPARTMENT OF HEALTH
Division of Epidemiology, Disease Surveillance and Investigation
899 N. Capitol Street, NE, Suite 580
Washington, D.C. 20002
(202) 442-9371 Fax (202) 442-8060
www.doh.dc.gov

What are noroviruses?

Norovirus is a virus that causes gastroenteritis, or vomiting and diarrhea, in people.

What are the symptoms of illness caused by noroviruses?

Norovirus illness usually begins 24 - 48 hours after exposure, but can appear as early as 10 hours after exposure. Symptoms usually include nausea, vomiting, diarrhea, and stomach cramping. Sometimes people have a low-grade fever, chills, headache, muscle aches, and a general sense of tiredness. The illness is usually brief, with symptoms lasting only 1 or 2 days.

How serious is norovirus disease?

Norovirus disease is usually not serious, but people may feel very sick. Most people get better within 1 or 2 days, and have no long-term health effects from the illness. Sometimes people are unable to drink enough liquids to replace what they lose from vomiting and diarrhea, and they can become dehydrated and need to see a doctor. This problem usually occurs only among the very young, the elderly, and persons with weakened immune systems.

How is norovirus spread?

Noroviruses are very contagious and spread easily from person to person. The virus is found in the stool and vomit of infected people. People can become infected in several ways, including eating food or drinking liquids that are contaminated by infected food handlers, touching surfaces or objects contaminated with norovirus and then touching their mouth before handwashing, or having direct contact with another person who is infected and then touching their mouth before handwashing.

How long are people contagious?

People infected with norovirus are contagious from the moment they begin feeling ill to at least 3 days after recovery. Some people may be contagious for as long as 2 weeks after recovery. Therefore, good handwashing is important. Persons infected with norovirus should not prepare food while they have symptoms and for 3 days

after they recover. Infected people do not become long-term carriers of norovirus.

Who gets norovirus infection?

Anyone can become infected with these viruses.

What treatment is available for people with norovirus infection?

Currently, there is no specific medication or vaccine for norovirus. Norovirus infection cannot be treated with antibiotics. By drinking fluids, such as juice or water, people can reduce their chance of becoming dehydrated.

Do infected people need to be excluded from school, work or daycare?

Since the virus is passed in vomit and stool, children should not go to daycare or school while they have diarrhea or vomiting. Once illness ends, children can return to daycare, but handwashing must be strictly monitored. Persons who work in nursing homes, take care of patients, or handle food should stay out of work until at least three days after symptoms end.

Can norovirus infections be prevented?

You can decrease your chance of becoming infected with noroviruses by:

- Frequent handwashing with warm water and soap
- Promptly disinfecting contaminated surfaces with household chlorine bleach-based cleaners
- Washing soiled clothing and linens
- Avoiding food or water from sources that may be contaminated
- Cooking oysters completely to kill the virus

Want more information?

Information about Norovirus and other related health topics can be found at the website www.cdc.gov. The DC Department of Health promotes the health and safety of the District residents. For additional information, please call (202) 442-9371.

